Quality assessment							Number of patients		Effect			
Number of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Psychoeducation	any control	Relative (95% CI)	Absolute (95% Cl)	Quality	Importance
Carer health and well-being (depression) – follow-up (follow up: mean 4 weeks)												
1	randomised trials	serious ¹	not serious	serious ²	very serious	none	40	-	-	SMD 0.84 fewer (1.31 fewer to 0.36 fewer)		CRITICAL
Coror boolth and w	ell-being (burnout)	follow up (fol										1

Quality assessment							Number of patients		Effect			
Number of studies	Study design	Risk of bias	Inconsistency	Indirectness	Imprecision	Other considerations	Psychoeducation	any control	Relative (95% CI)	Absolute (95% Cl)	Quality	Importance
1	randomised trials	serious ¹	not serious	serious ²	very serious	none	45			SMD 0.35 fewer (0.77 fewer to 0.06 more)		CRITICAL

Crucial limitation for one criterion or some limitations for multiple criteria sufficient to lower ones confidence in the estimate of effect Population not family carers of people with learning disabilities with no mental health problems. Optimal information size not met; small, single study 1.

2. 3.

© National Guideline Alliance, 2016