



## Mitochondrial DNA Maintenance Defects Overview

Ayman W El-Hattab, MD, FAAP, FACMG,<sup>1</sup> William J Craigen, MD, PhD, FACMG,<sup>2</sup> Lee-Jun C Wong, PhD, FACMG,<sup>2</sup> and Fernando Scaglia, MD, FAAP, FACMG<sup>2</sup>

Created: March 8, 2018.

### Summary

This overview focuses on the clinical features and molecular genetics of mitochondrial DNA (mtDNA) maintenance defects.

The goals of this overview are the following.

#### Goal 1

Describe the pathomechanism of mtDNA maintenance defects.

#### Goal 2

Review the genetic causes of mtDNA maintenance defects.

#### Goal 3

Describe the clinical characteristics of mtDNA maintenance defects.

#### Goal 4

Provide clinical and laboratory evaluation strategies to facilitate the diagnosis of a mtDNA maintenance defect and to establish a genetic cause in a proband (when possible).

#### Goal 5

Inform genetic counseling for mtDNA maintenance defects.

#### Goal 6

Summarize current management recommendations for individuals with mtDNA maintenance defects.

**Author Affiliations:** 1 Associate Professor, Department of Clinical Sciences, College of Medicine, University of Sharjah, Sharjah, United Arab Emirates; Email: elhattabaw@yahoo.com. 2 Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, Texas; Email: wcraigen@bcm.edu; Email: ljwong@bcm.edu; Email: fscaglia@bcm.edu.

## 1. Mitochondrial DNA Maintenance Defects

The maintenance of mtDNA is essential to the functioning of the mitochondria and, thus, to meeting the energy needs of all cells. The maintenance of mtDNA requires proteins essential for mtDNA synthesis, for maintenance of the mitochondrial nucleotide pool, and for mediating mitochondrial fusion [El-Hattab et al 2017].

Mitochondrial DNA is synthesized continuously and is not regulated by the cell cycle. The enzymes that synthesize mtDNA require a balanced supply of intramitochondrial nucleotides. These are supplied through mitochondrial nucleotide salvage pathways and the import of nucleotides from the cytosol via specific transporters. To function properly in mtDNA synthesis the quantities of these enzymes need to be perfectly balanced, a phenomenon achieved – in part – by the exchange of content between mitochondria through the process of mitochondrial fission and fusion.

The proteins known to be required for mtDNA synthesis are encoded by nuclear genes (i.e., genes found in the nucleus of cells). When pathogenic variants disrupt the function of any one of the proteins encoded by these genes, mtDNA synthesis is impaired, resulting in either quantitative defects in mtDNA (mtDNA depletion) or qualitative defects in mtDNA (multiple mtDNA deletions). These defects in mtDNA maintenance result in energy deficiency within cells. Cellular energy production insufficient to meet the needs of a given organ results in organ dysfunction (see Figure 1).

When first identified, defects in mtDNA maintenance were viewed as two clinically distinct groups of disorders:


- Mitochondrial DNA depletion syndromes that typically present during infancy and are characterized by severe disease manifestations and shortened life expectancy; and
- Multiple mtDNA deletion syndromes that typically present in adulthood and are characterized by milder disease manifestations including progressive external ophthalmoplegia (CPEO) and myopathy.

However, with the current understanding that both mtDNA depletion and multiple mtDNA deletions result from failure of proper mtDNA maintenance, it has become evident that these two groups of disorders represent the ends of a phenotypic continuum. The term "mtDNA maintenance defects" is used to represent the broad disease spectrum that encompasses both presentations as well as those that are intermediate.

## 2. Causes of mtDNA Maintenance Defects

To date, pathogenic variants in 20 nuclear genes are known to be associated with mtDNA maintenance defects. These genes and their primary presenting features are organized in Table 1 by the category of defect: mtDNA synthesis, mitochondrial nucleotide salvage pathway, cytosolic nucleotide metabolism, mitochondrial nucleotide import, and mitochondrial fusion.

Note: Disorders of mtDNA are not the subject of this overview (see [Primary Mitochondrial Disorders Overview](#)).



**Figure 1.** A diagram showing the proteins that are involved in mtDNA maintenance and known to be associated with mtDNA maintenance defects

**Enzymes of the mitochondrial nucleotide salvage pathway.** Thymidine kinase 2 (TK2; encoded by *TK2*) and deoxyguanosine kinase (DGK; encoded by *DGUOK*) convert the deoxyribonucleosides to deoxyribonucleotide monophosphates. Nucleotide monophosphate

kinase (NMPK) converts deoxyribonucleotide monophosphates to deoxyribonucleotide diphosphates. Nucleotide diphosphate kinase (NDPK) converts deoxyribonucleotide diphosphates to deoxyribonucleotide triphosphates. NDPK forms a complex with both succinyl-CoA ligase (SUCL) (composed of an alpha subunit encoded by *SUCLG1* and a beta subunit encoded by either *SUCLA2* or *SUCLG2*) and gamma-aminobutyrate transaminase (GABAT), encoded by *ABAT*.

**Enzymes of cytosolic nucleotide metabolism.** Ribonucleotide reductase (RNR) composed of two catalytic subunits and two small subunits (either R2 or p53-inducible small RNR subunit [p53R2; encoded by *RRM2B*]) converts ribonucleotide diphosphates to deoxyribonucleotide diphosphates. Thymidine phosphorylase (TP; encoded by *TYMP*) converts thymidine to thymine.

**Proteins involved in mitochondrial nucleotide transport.** Adenine nucleotide translocator 1 (ANT1; encoded by *SLC25A4*), acylglycerol kinase (AGK; encoded by *AGK*), and MPV17 (encoded by *MPV17*)

**Enzymes involved in mtDNA replication.** Twinkle (encoded by *TWNK*), a DNA helicase that separates the DNA strands (DNA polymerase  $\gamma$  [pol  $\gamma$ ], consisting of a catalytic subunit encoded by *POLG*, and two accessory subunits encoded by *POLG2*) requires an RNA primer to initiate DNA replication. Mitochondrial transcription factor A (TFAM; encoded by *TFAM*), is required for the generation of an RNA primer. Nucleases removing RNA primers and flap intermediates are RNase H1 (encoded by *RNASEH1*), DNA helicase/nuclease 2 (DNA2; encoded by *DNA2*), and mitochondrial genome maintenance exonuclease 1 (MGME1; encoded by *MGME1*).

**Proteins involved in mitochondrial fusion.** Mitofusin 2 (MFN2; encoded by *MFN2*), dynamin-related GTPase OPA1 (encoded by *OPA1*), and F-box and leucine-rich repeat 4 (FBXL4; encoded by *FBXL4*)

**Table 1.** Categories of mtDNA Maintenance Defects: Genes and Primary Presenting Features

Category of Defect	Gene	Primary Presenting Features							
		Encephalo-hepatopathy	Encephalo-myopathy	Encephalo-neuropathy	Neurogastro-intestinal Encephalopathy	Myopathy	Ophthal-moplegia	Optic Atrophy	Neuropathy
Mitochondrial DNA synthesis	<i>POLG</i>	X	X	X	X	X	X		
	<i>POLG2</i>					X			
	<i>TWNK</i>	X		X			X		
	<i>TFAM</i>	X							
	<i>RNASEH1</i>		X						
	<i>MGME1</i>					X			
	<i>DNA2</i>					X			
Mitochondrial nucleotide salvage pathway	<i>TK2</i>					X	X		
	<i>DGUOK</i>	X				X			
	<i>SUCLA2</i>		X						
	<i>SUCLG1</i>		X						
Cytosolic nucleotide metabolism	<i>ABAT</i>		X						
	<i>TYMP</i>				X				
Mitochondrial nucleotide import	<i>RRM2B</i>		X		X		X		
	<i>SLC25A4</i>					X	X		
Mitochondrial fusion	<i>AGK</i>					X			
	<i>MPV17</i>	X							
	<i>OPA1</i>		X	X				X	
Mitochondrial fusion	<i>MFN2</i>							X	X
	<i>FBXL4</i>		X						

### 3. Clinical Characteristics of mtDNA Maintenance Defects

Mitochondrial DNA maintenance defects are characterized by mtDNA depletion and/or multiple mtDNA deletions in mitochondria of cells of affected organs. The organs/tissues affected most often are the brain, liver, skeletal muscle, peripheral nerves, and gastrointestinal tract. Depending on the organ(s) predominantly affected, these disorders can be classified into groups associated mainly with encephalohepatopathy (see Table 2a), encephalomyopathy (Table 2b), encephaloneuropathy (Table 2c), neurogastrointestinal encephalopathy (Table 2d), myopathy (Table 2e), ophthalmoplegia (Table 2f), optic atrophy (Table 2g), or neuropathy (Table 2h).

#### Mitochondrial DNA Maintenance Defects Presenting with Encephalohepatopathy

Mitochondrial DNA maintenance defects manifesting as encephalohepatopathy (hepatocerebral) are typically associated with mtDNA depletion and generally present in neonates or infants with neurologic manifestations (including developmental delay and epilepsy), and with liver dysfunction and failure. Other common manifestations include growth failure, lactic acidosis, and hypoglycemia.

**Table 2a.** Mitochondrial DNA Maintenance Defects Presenting with Encephalohepatopathy

Gene	Disorder/Phenotype	MOI	mtDNA Maintenance Defect	Usual Age of Onset	Common Manifestations in Addition to Liver Dysfunction/Failure
<i>DGUOK</i>	Deoxyguanosine kinase deficiency	AR	Depletion	Neonatal	<ul style="list-style-type: none"> <li>• DD</li> <li>• Hypotonia</li> <li>• Nystagmus</li> <li>• Lactic acidosis</li> </ul>
<i>MPV17</i>	Hepatocerebral mtDNA depletion syndrome	AR	Depletion	Neonatal or infancy	<ul style="list-style-type: none"> <li>• DD</li> <li>• Hypotonia</li> <li>• Failure to thrive</li> <li>• Hearing impairment</li> <li>• Lactic acidosis</li> </ul>
<i>POLG</i>	Alpers-Huttenlocher syndrome	AR	Depletion	Early childhood	<ul style="list-style-type: none"> <li>• DD</li> <li>• Psychomotor regression</li> <li>• Epilepsy</li> <li>• Hearing impairment</li> </ul>
<i>TFAM</i>	Encephalohepatopathy (OMIM 617156)	AR	Depletion	Neonatal	<ul style="list-style-type: none"> <li>• IUGR</li> <li>• Hypoglycemia</li> </ul>
<i>TWNK</i>	Encephalohepatopathy (OMIM 271245)	AR	Depletion	Neonatal or infancy	<ul style="list-style-type: none"> <li>• DD</li> <li>• Hypotonia</li> <li>• Lactic acidosis</li> </ul>

AR = autosomal recessive; DD = developmental delay; IUGR = intrauterine growth restriction; MOI = mode of inheritance; mtDNA = mitochondrial DNA

#### Mitochondrial DNA Maintenance Defects Presenting with Encephalomyopathy

The majority of encephalomyopathic mtDNA maintenance defects are associated with mtDNA depletion and are early-onset diseases with an infantile presentation. The two disorders, however, that are usually associated with multiple mtDNA deletions rather than depletion are adult-onset diseases: *POLG*-related myoclonic epilepsy-myopathy-sensory ataxia and *RNASEH1*-related encephalomyopathy.

**Table 2b.** Mitochondrial DNA Maintenance Defects Presenting with Encephalomyopathy

Gene	Disorder	MOI	mtDNA Maintenance Defect	Usual Age of Onset	Common Manifestations in Addition to Muscle Weakness
<i>ABAT</i>	Encephalomyopathy w/elevated GABA (OMIM 613163)	AR	Depletion	Infancy	<ul style="list-style-type: none"> <li>• DD</li> <li>• Hypotonia</li> <li>• Epilepsy</li> <li>• ↑ GABA in plasma, urine, &amp; CSF</li> </ul>
<i>FBXL4</i>	Encephalomyopathic mtDNA depletion syndrome	AR	Depletion	Neonatal or infancy	<ul style="list-style-type: none"> <li>• DD</li> <li>• Hypotonia</li> <li>• Epilepsy</li> <li>• Hearing impairment</li> <li>• Lactic acidosis</li> </ul>
<i>OPA1</i>	Encephalomyopathy (OMIM 616896)	AR	Depletion	Infancy	<ul style="list-style-type: none"> <li>• DD</li> <li>• HCM</li> <li>• Optic atrophy</li> </ul>
<i>POLG</i>	Myoclonic epilepsy-myopathy-sensory ataxia	AR	Multiple deletions	Early adulthood	<ul style="list-style-type: none"> <li>• Epilepsy</li> <li>• Ataxia</li> </ul>
<i>RNASEH1</i>	Encephalomyopathy (OMIM 616479)	AR	Depletion & multiple deletions	Early adulthood	<ul style="list-style-type: none"> <li>• Ophthalmoplegia</li> <li>• Ptosis</li> <li>• Ataxia</li> </ul>
<i>RRM2B</i>	Encephalomyopathy w/renal tubulopathy	AR	Depletion	Neonatal or infancy	<ul style="list-style-type: none"> <li>• DD</li> <li>• Hypotonia</li> <li>• GI dysmotility</li> <li>• Renal tubulopathy</li> </ul>
<i>SUCLA2</i>	Mitochondrial DNA depletion syndrome, encephalomyopathic form w/methylmalonic aciduria	AR	Depletion	Infancy or early childhood	<ul style="list-style-type: none"> <li>• DD</li> <li>• Hypotonia</li> <li>• Dystonia</li> <li>• Hearing impairment</li> <li>• ↑ methylmalonic acid</li> </ul>
<i>SUCLG1</i>	Mitochondrial DNA depletion syndrome, encephalomyopathic form w/methylmalonic aciduria	AR	Depletion	Neonatal or infancy	<ul style="list-style-type: none"> <li>• DD</li> <li>• Hypotonia</li> <li>• Hearing impairment</li> <li>• ↑ methylmalonic acid</li> </ul>

AR = autosomal recessive; CSF = cerebrospinal fluid; DD = developmental delay; GI = gastrointestinal; MOI = mode of inheritance; mtDNA = mitochondrial DNA

## Mitochondrial DNA Maintenance Defects Presenting with Encephaloneuropathy

Mitochondrial DNA maintenance defects exhibiting encephaloneuropathy can be associated with mtDNA depletion or multiple mtDNA deletions, and are characterized by manifestations related to the central and peripheral nervous systems.

**Table 2c.** Mitochondrial DNA Maintenance Defects Presenting with Encephaloneuropathy

Gene	Disorder	MOI	mtDNA Maintenance Defect	Usual Age of Onset	Common Manifestations in Addition to Peripheral Neuropathy & Ataxia
<i>OPA1</i>	Behr syndrome (OMIM 210000)	AR	NA	Infancy or early childhood	<ul style="list-style-type: none"> <li>Vision impairment</li> <li>Optic nerve pallor</li> </ul>
<i>POLG</i>	Ataxia neuropathy spectrum disorders	AR	Multiple deletions	Early adulthood	Epilepsy
<i>TWINK</i>	Infantile-onset spinocerebellar ataxia	AR	Depletion	2nd year of life	<ul style="list-style-type: none"> <li>Hypotonia</li> <li>Hearing impairment</li> </ul>

AR = autosomal recessive; MOI = mode of inheritance; mtDNA = mitochondrial DNA

## Mitochondrial DNA Maintenance Defects Presenting with Neurogastrointestinal Encephalopathy

Mitochondrial neurogastrointestinal encephalopathy (MNGIE) is characterized by a variable age of onset (generally in the 2nd decade) and progressive gastrointestinal dysmotility, peripheral neuropathy, and leukoencephalopathy. The gastrointestinal manifestations of the disease may mimic anorexia nervosa. MNGIE is most commonly caused by biallelic pathogenic variants in *TYMP*, the gene encoding thymidine phosphorylase; however, biallelic pathogenic variants in *POLG* or *RRM2B* also cause this disorder.

**Table 2d.** Mitochondrial DNA Maintenance Defects Presenting with Neurogastrointestinal Encephalopathy

Gene	Disorder	MOI	mtDNA Maintenance Defect	Usual Age of Onset	Common Manifestations
<i>TYMP</i>	MNGIE type 1	AR	Depletion & multiple deletions	Adolescence or early adulthood	<ul style="list-style-type: none"> <li>GI dysmotility</li> <li>Cachexia</li> </ul>
<i>POLG</i>	MNGIE type 4B	AR	Depletion & multiple deletions	Infancy or childhood	<ul style="list-style-type: none"> <li>Peripheral neuropathy</li> <li>Ophthalmoplegia</li> </ul>
<i>RRM2B</i>	MNGIE type 8B	AR	Depletion	Early adulthood	<ul style="list-style-type: none"> <li>Muscle weakness</li> <li>Leukoencephalopathy<sup>1</sup></li> </ul>

AR = autosomal recessive ; GI = gastrointestinal; MOI = mode of inheritance; mtDNA = mitochondrial DNA

1. Note: Leukoencephalopathy is not present in *POLG*-related neurogastrointestinal encephalopathy.

## Mitochondrial DNA Maintenance Defects Presenting with Myopathy

Myopathic mtDNA maintenance defects include a group of diseases that vary in their age of onset. Skeletal muscles are the main system involved in all of them. Cardiomyopathy can occur in some of these disorders.

**Table 2e.** Mitochondrial DNA Maintenance Defects Presenting with Myopathy

Gene	Disorder	MOI	mtDNA Maintenance Defect	Usual Age of Onset	Common Clinical Manifestations in Addition to Muscle Weakness
<i>AGK</i>	Sengers syndrome (OMIM 212350)	AR	Depletion	Neonatal period	<ul style="list-style-type: none"> <li>Hypotonia</li> <li>Hypertrophic cardiomyopathy</li> <li>Cataracts</li> </ul>
<i>DGUOK</i>	Myopathy	AR	Multiple deletions	Early or mid-adulthood	<ul style="list-style-type: none"> <li>Ptosis</li> <li>Ophthalmoplegia</li> </ul>

Table 2e. continued from previous page.

Gene	Disorder	MOI	mtDNA Maintenance Defect	Usual Age of Onset	Common Clinical Manifestations in Addition to Muscle Weakness
<i>DNA2</i>	Myopathy (OMIM 615156)	AD	Multiple deletions	Childhood or early adulthood	<ul style="list-style-type: none"> <li>• Ptosis</li> <li>• Ophthalmoplegia</li> </ul>
<i>MGME1</i>	Myopathy (OMIM 615084)	AR	Depletion & multiple deletions	Childhood or early adulthood	<ul style="list-style-type: none"> <li>• Ptosis</li> <li>• Ophthalmoplegia</li> </ul>
<i>POLG2</i>	Myopathy (OMIM 610131)	AD	Multiple deletions	Infancy to adulthood	<ul style="list-style-type: none"> <li>• Ptosis</li> <li>• Ophthalmoplegia</li> </ul>
<i>SLC25A4</i>	Cardiomyopathy (OMIM 615418)	AR	Multiple deletions	Childhood	<ul style="list-style-type: none"> <li>• Exercise intolerance / easy fatigability</li> <li>• Hypertrophic cardiomyopathy</li> </ul>
	Cardiomyopathy (OMIM 617184)	AD	Depletion	Birth	<ul style="list-style-type: none"> <li>• Hypotonia</li> <li>• Hypertrophic cardiomyopathy</li> </ul>
<i>TK2</i>	Mitochondrial DNA depletion syndrome	AR	Depletion	Infancy or childhood	<ul style="list-style-type: none"> <li>• Hypotonia</li> <li>• Loss of acquired motor skills</li> </ul>

AD = autosomal dominant; AR = autosomal recessive; MOI = mode of inheritance; mtDNA = mitochondrial DNA

## Mitochondrial DNA Maintenance Defects Presenting with Ophthalmoplegia

Mitochondrial DNA maintenance defects that cause ophthalmoplegia are associated with multiple DNA deletions and are characterized by progressive weakness of the extraocular eye muscles resulting in ptosis (drooping of the eyelids) and ophthalmoplegia (paralysis of the extraocular muscles causing limitation in horizontal and vertical eye movements).

Although these are typically diseases of adulthood, earlier onset can be seen in the recessively inherited diseases. Although ophthalmoplegia and ptosis are consistent, and are the main manifestations in these diseases, a more generalized myopathy (sometimes mild) can be observed in some affected individuals.

Table 2f. Mitochondrial DNA Maintenance Defects Presenting with Ophthalmoplegia

Gene	Disorder	MOI	mtDNA Maintenance Defect	Usual Age of Onset	Common Clinical Manifestations in Addition to Ptosis & Ophthalmoplegia
<i>POLG</i>	Progressive external ophthalmoplegia	AR	Multiple deletions	Adolescence or young adulthood	Easy fatigability / exercise intolerance
		AD	Multiple deletions	Adulthood	Easy fatigability / exercise intolerance
<i>RRM2B</i>	Chronic progressive external ophthalmoplegia	AR	Multiple deletions	Childhood	<ul style="list-style-type: none"> <li>• Muscle weakness</li> <li>• Bulbar dysfunction</li> </ul>
		AD	Multiple deletions	Adulthood	<ul style="list-style-type: none"> <li>• Ataxia</li> <li>• Muscle weakness</li> <li>• Bulbar dysfunction</li> </ul>
<i>SLC25A4</i>	Progressive external ophthalmoplegia (OMIM 609283)	AD	Multiple deletions	Adulthood	Easy fatigability / exercise intolerance
<i>TK2</i>	Progressive external ophthalmoplegia (OMIM 617069)	AR	Multiple deletions	Adulthood	Muscle weakness



Table 2f. continued from previous page.

Gene	Disorder	MOI	mtDNA Maintenance Defect	Usual Age of Onset	Common Clinical Manifestations in Addition to Ptosis & Ophthalmoplegia
<i>TWINK</i>	Progressive external ophthalmoplegia (OMIM 609286)	AD	Multiple deletions	Early adulthood	Easy fatigability / exercise intolerance

AD = autosomal dominant; AR = autosomal recessive; MOI = mode of inheritance; mtDNA = mitochondrial DNA

## Mitochondrial DNA Maintenance Defects Presenting with Optic Atrophy

Table 2g. Mitochondrial DNA Maintenance Defects Presenting with Optic Atrophy

Gene	Disorder	MOI	mtDNA Maintenance Defect	Usual Age of Onset	Common Clinical Manifestations
<i>OPA1</i>	Optic atrophy type 1	AD	Multiple deletions	Childhood	Vision impairment
<i>MFN2</i>	Optic atrophy	AD	Multiple deletions	Early childhood	<ul style="list-style-type: none"> <li>• Vision impairment</li> <li>• Optic nerve pallor</li> <li>• Peripheral neuropathy</li> <li>• Muscle weakness</li> </ul>

AD = autosomal dominant; MOI = mode of inheritance

## Mitochondrial DNA Maintenance Defects Presenting with Neuropathy

Table 2h. Mitochondrial DNA Maintenance Defects Presenting with Neuropathy

Gene	Disorder	MOI	mtDNA Maintenance Defect	Usual Age of Onset	Common Clinical Manifestations
<i>MFN2</i>	Charcot-Marie-Tooth neuropathy type 2a	AD	NA	Childhood or early adulthood	Peripheral neuropathy

AD = autosomal dominant; MOI = mode of inheritance; mtDNA = mitochondrial DNA; NA= not available

## 4. Evaluation Strategies to Diagnose mtDNA Maintenance Defects and to Establish a Genetic Cause in a Proband

Establishing a specific genetic cause of a mtDNA maintenance defect can aid in discussion of prognosis (which is beyond the scope of this *GeneReview*) and in genetic counseling. See Genetic Counseling.

Establishing the specific genetic cause of a mtDNA maintenance defect usually requires a medical history, physical and neurologic examination, laboratory testing including routine studies and specialized biochemical genetic studies, imaging studies such as brain MRI, echocardiogram, abdominal ultrasound examination, family history, and genomic/genetic testing.

### Clinical and Laboratory Findings

The diagnosis of a mtDNA maintenance defect is suspected based on the involved organs, age of onset, and results of commonly available laboratory tests (e.g., presence of lactic acidemia or methylmalonic aciduria).

Biopsies of affected tissues typically show mtDNA depletion and/or multiple mtDNA deletions as well as decreased activity of multiple electron transport complexes (ETC). However, because tissue biopsies are often invasive procedures, molecular genetic testing of leukocyte DNA is typically performed first to determine if the diagnosis of a mtDNA maintenance defect can be established. When molecular genetic test results are equivocal

or fail to confirm the diagnosis of a mtDNA maintenance defect, tissue samples can be obtained to assay for mtDNA depletion and/or multiple mtDNA deletions and to assess ETC activity.

## Family History

A three-generation family history should be obtained, with attention to relatives with signs and symptoms that could be related to a mtDNA maintenance defect and documentation of relevant findings through direct physical examination, or review of medical records, including results of molecular genetic testing.

## Molecular Genetic Testing

Approaches include gene-targeted testing (multigene panel or single-gene testing) or comprehensive genomic testing (exome sequencing). Gene-targeted testing requires the clinician to hypothesize which gene(s) are likely involved, whereas genomic testing may not. Options for testing include the following:

- **Serial single-gene testing** can be considered if clinical findings and/or family history indicate that mutation of a particular gene is most likely (see Tables 2a-2h). Sequence analysis of the gene of interest can detect small intragenic deletions/insertions and missense, nonsense, and splice site variants; typically, exon or whole-gene deletions/duplications are not detected by sequence analysis, therefore, deletion/duplication analysis should also be performed to detect intragenic deletions or duplications.
- **A multigene panel** that includes some or all of the mtDNA maintenance genes (Table 1) is most likely to identify the genetic cause of the condition while limiting identification of variants of uncertain significance and pathogenic variants in genes that do not explain the underlying phenotype. Note: (1) The genes included in the panel and the diagnostic sensitivity of the testing used for each gene vary by laboratory and are likely to change over time. (2) Some multigene panels may not include all the genes discussed in this *GeneReview*. (3) In some laboratories, panel options may include a custom laboratory-designed panel and/or custom phenotype-focused exome analysis that includes genes specified by the clinician. (4) Methods used in a panel may include sequence analysis, deletion/duplication analysis, and/or other non-sequencing-based tests.

For an introduction to multigene panels click [here](#). More detailed information for clinicians ordering genetic tests can be found [here](#).

- **Comprehensive genomic testing** (which does not require the clinician to determine which gene[s] are likely involved) can be considered. **Exome sequencing** is most commonly used; **genome sequencing** is also possible.

For an introduction to comprehensive genomic testing click [here](#). More detailed information for clinicians ordering genomic testing can be found [here](#).

## 5. Genetic Counseling

*Genetic counseling is the process of providing individuals and families with information on the nature, mode(s) of inheritance, and implications of genetic disorders to help them make informed medical and personal decisions. The following section deals with genetic risk assessment and the use of family history and genetic testing to clarify genetic status for family members; it is not meant to address all personal, cultural, or ethical issues that may arise or to substitute for consultation with a genetics professional. —ED.*

## Mode of Inheritance

Mitochondrial DNA maintenance defects can be inherited in an autosomal recessive or autosomal dominant manner.

## Autosomal Recessive Inheritance – Risk to Family Members

### Parents of a proband

- The parents of an affected child are obligate heterozygotes (i.e., carriers of one mtDNA maintenance defect-related pathogenic variant).
- Heterozygotes (carriers) are asymptomatic and are not at risk of developing the disorder.

### Sibs of a proband

- At conception, each sib of an affected individual has a 25% chance of being affected, a 50% chance of being an asymptomatic carrier, and a 25% chance of being unaffected and not a carrier.
- Heterozygotes (carriers) are asymptomatic and are not at risk of developing the disorder.

**Offspring of a proband.** Offspring of an individual with a mtDNA maintenance defect are obligate heterozygotes (carriers) for a pathogenic variant.

**Other family members.** Each sib of the proband's parents is at a 50% risk of being a carrier of a mtDNA maintenance defect-related pathogenic variant.

**Carrier detection.** Carrier testing for at-risk relatives requires prior identification of the pathogenic variants in the family.

## Autosomal Dominant Inheritance – Risk to Family Members

### Parents of a proband

- Most individuals diagnosed with a mtDNA maintenance defect have an affected parent.
- Some individuals diagnosed with a mtDNA maintenance defect have the disorder as the result of a *de novo* pathogenic variant.
- Molecular genetic testing is recommended for the parents of a proband with an apparent *de novo* pathogenic variant.
- If the pathogenic variant found in the proband cannot be detected in leukocyte DNA of either parent, possible explanations include a *de novo* pathogenic variant in the proband or germline mosaicism in a parent. Though theoretically possible, no instances of germline mosaicism have been reported to date.
- The family history of some individuals diagnosed with a mtDNA maintenance defect may appear to be negative because of failure to recognize the disorder in family members, early death of the parent before the onset of symptoms, or late onset of the disease in the affected parent. Therefore, an apparently negative family history is not definitive unless appropriate clinical evaluation and/or molecular genetic testing has been performed for the parents of the proband.

### Sibs of a proband

- The risk to sibs of the proband depends on the genetic status of the proband's parents.
- If a parent of the proband is affected and/or is heterozygous for the pathogenic variant identified in the proband, the risk to sibs is 50%.
- If the pathogenic variant found in the proband cannot be detected in the leukocyte DNA of either parent, the recurrence risk to sibs is estimated to be 1% because of the theoretic possibility of parental germline mosaicism [Rahbari et al 2016].
- If the parents have not been tested for the mtDNA maintenance defect-related pathogenic variant but are clinically unaffected, the risk to sibs of a proband appears to be low.

**Offspring of a proband.** Each child of an individual with a mtDNA maintenance defect has a 50% chance of inheriting the pathogenic variant.

**Other family members.** The risk to other family members depends on the status of the proband's parents: if a parent has the mtDNA maintenance defect-related pathogenic variant, the parent's family members may be at risk.

## Prenatal Testing and Preimplantation Genetic Testing

Once the mtDNA maintenance defect-related pathogenic variant(s) have been identified in an affected family member, prenatal and preimplantation genetic testing are possible.

Differences in perspective may exist among medical professionals and within families regarding the use of prenatal testing. While most centers would consider use of prenatal testing to be a personal decision, discussion of these issues may be helpful.

## 6. Management of Individuals with mtDNA Maintenance Defects

Most mtDNA maintenance defects affect multiple organs; therefore, affected individuals need comprehensive evaluations to assess the degree of involvement of different organs. Management should also involve a multidisciplinary team to provide clinical care for these multiorgan diseases.

### Assessment of Disease Extent

For individuals with chronic disease, Table 3 summarizes evaluations that are recommended if they have not already been completed:

**Table 3.** Recommended Evaluations Following Initial Diagnosis in a Proband with a mtDNA Maintenance Defect Resulting in Chronic Disease

System/ Concern	Evaluation	Comment
<b>Eyes</b>	Ophthalmologic eval (See <a href="#">POLG-Related Disorders</a> .)	To assess for optic atrophy, ptosis, ophthalmoplegia, & nystagmus
<b>ENT/Mouth</b>	Hearing eval (See <a href="#">POLG-Related Disorders</a> .)	
<b>Cardiovascular</b>	<ul style="list-style-type: none"> <li>Echocardiogram</li> <li>Electrocardiogram</li> </ul>	For persons w/myopathy
<b>Respiratory</b>	<ul style="list-style-type: none"> <li>Venous blood gases</li> <li>Pulse oximetry &amp; pulmonary function tests</li> <li>Polysomnography</li> </ul>	To identify respiratory insufficiency in persons w/ myopathy
<b>Gastrointestinal</b>	<ul style="list-style-type: none"> <li>Liver function test (transaminases, albumin, coagulation profile)</li> <li>Liver ultrasound</li> </ul>	For persons w/hepatopathy
	For MNGIE disease (see <a href="#">Mitochondrial Neurogastrointestinal Encephalopathy Disease</a> ): <ul style="list-style-type: none"> <li>Consultation w/gastroenterologist</li> <li>Depending on manifestations: abdominal films, abdominal CT, upper GI contrast radiography, esophagogastroduodenoscopy, sigmoidoscopy, liquid phase scintigraphy, antroduodenal manometry</li> </ul>	To evaluate for gastrointestinal dysmotility
<b>Feeding</b>	<ul style="list-style-type: none"> <li>Swallowing assessment</li> <li>Nutritional eval</li> </ul>	In individuals w/feeding difficulty & growth failure <sup>1</sup>
<b>Renal</b>	<ul style="list-style-type: none"> <li>Urinalysis</li> <li>Urine amino acids, calcium, phosphate, &amp; protein</li> </ul>	To evaluate for tubulopathy

Table 3. continued from previous page.

System/ Concern	Evaluation	Comment
<b>Neurologic</b>	<ul style="list-style-type: none"> <li>• Comprehensive neurologic exam</li> <li>• Brain MRI &amp; MRS</li> <li>• Nerve conduction studies &amp; electromyography (if neuropathy is suspected)</li> <li>• Electroencephalography (if seizures are suspected)</li> </ul>	For persons w/neurologic manifestations
<b>Musculoskeletal</b>	Referral to rehabilitation specialist	Evaluate gait, weakness, safety, activities of daily living
<b>Genetics &amp; metabolic</b>	<ul style="list-style-type: none"> <li>• Consultation w/clinical geneticist &amp;/or genetic counselor</li> <li>• Lactate level to evaluate for lactic acidosis</li> <li>• Glucose level to evaluate for hypoglycemia</li> </ul>	

1. El-Hattab et al [2017]

## Treatment of Manifestations

Currently there is no clinical therapy to treat the primary defect in affected individuals. Management, which is primarily supportive, is outlined in Table 4. Some specific considerations:

- As exogenous thymidine phosphorylase can improve outcome in MNGIE resulting from thymidine phosphorylase deficiency, experimental therapy for MNGIE includes both bone marrow and liver transplantation.
- Nucleoside therapy has been considered in TK2 deficiency.
- Affected individuals may be at increased risk for acidosis and hypoglycemia during illness and surgery and protocols to prevent prolonged fasting should be provided.
- Certain medications and anesthetic agents should be avoided; see [Primary Mitochondrial Disorders Overview](#).

**Table 4.** Treatment of Manifestations in a Proband with a mtDNA Maintenance Defect Resulting in Chronic Disease

Manifestation/ Concern	Treatment
<b>Ptosis</b>	Ptosis blepharoplasty
<b>Sensorineural hearing loss</b>	Hearing aids & cochlear implantation (See <a href="#">POLG-Related Disorders</a> , <a href="#">RRM2B Mitochondrial DNA Maintenance Defects</a> , <a href="#">Hereditary Hearing Loss and Deafness Overview</a> .)
<b>Cardiomyopathy, hypertrophic or dilated</b>	<ul style="list-style-type: none"> <li>• Referral to cardiologist</li> <li>• Standard treatment</li> </ul>
<b>Respiratory insufficiency</b>	<ul style="list-style-type: none"> <li>• Referral to pulmonologist &amp;/or sleep medicine physician</li> <li>• Aggressive antibiotic treatment of chest infections</li> <li>• Chest physiotherapy</li> <li>• Artificial ventilation including assisted nasal ventilation (CPAP or BiPAP) or intubation w/use of tracheostomy &amp; ventilator (See <a href="#">TK2-Related Mitochondrial DNA Depletion Syndrome</a>, <a href="#">Myopathic Form</a>.)</li> </ul>
<b>Liver failure</b>	<ul style="list-style-type: none"> <li>• Referral to hepatologist</li> <li>• Reduction in dietary protein</li> <li>• Correction of coagulopathy</li> <li>• Frequent or continuous feeding to prevent hypoglycemia</li> <li>• Consideration of liver transplant</li> </ul>

Table 4. continued from previous page.

Manifestation/ Concern	Treatment
<b>Gastrointestinal dysmotility</b>	<ul style="list-style-type: none"> <li>• Referral to gastroenterologist</li> <li>• Nutritional support</li> <li>• Total parenteral nutrition</li> <li>• Domperidone</li> <li>• Antibiotic therapy for intestinal bacterial overgrowth</li> <li>• Celiac plexus &amp; splanchnic nerve block (See <a href="#">Mitochondrial Neurogastrointestinal Encephalopathy Disease</a>.)</li> </ul>
<b>Failure to thrive &amp; feeding difficulties</b>	<ul style="list-style-type: none"> <li>• Nutritional support</li> <li>• Gastrostomy tube placement</li> </ul>
<b>Renal tubulopathy</b>	<ul style="list-style-type: none"> <li>• Referral to nephrologist</li> <li>• Correction of acidosis &amp; other metabolic derangements</li> </ul>
<b>Neuropathy</b>	<ul style="list-style-type: none"> <li>• Referral to neurologist</li> <li>• Amitriptyline, nortriptyline, &amp; gabapentin</li> </ul>
<b>Seizures</b>	<ul style="list-style-type: none"> <li>• Referral to neurologist</li> <li>• Standard ASM (Refractory epilepsy may require high doses &amp;/or use of multiple ASMs.)</li> </ul>
<b>Hypoglycemia</b>	<ul style="list-style-type: none"> <li>• Frequent feeding &amp; avoidance of fasting</li> <li>• Uncooked cornstarch</li> </ul>

ASM = anti-seizure medication

## Developmental Delay / Intellectual Disability Management Issues

The following information represents typical management recommendations for individuals with developmental delay / intellectual disability in the United States; standard recommendations may vary from country to country.

**Ages 0-3 years.** Referral to an early intervention program is recommended for access to occupational, physical, speech, and feeding therapy. In the US, early intervention is a federally funded program available in all states.

**Ages 3-5 years.** In the US, developmental preschool through the local public school district is recommended. Before placement, an evaluation is made to determine needed services and therapies and an individualized education plan (IEP) is developed.

### Ages 5-21 years

- In the US, an IEP based on the individual's level of function should be developed by the local public school district. Affected children are permitted to remain in the public school district until age 21.
- Discussion about transition plans including financial, vocation/employment, guardianship, and medical arrangements should begin at age 12 years. Developmental pediatricians can provide assistance with transition to adulthood.

**All ages.** Consultation with a developmental pediatrician is recommended to ensure the involvement of appropriate community, state, and educational agencies and to support parents in maximizing quality of life.

Consideration of private supportive therapies based on the affected individual's needs is recommended. Specific recommendations regarding type of therapy can be made by a developmental pediatrician.

In the US:

- Developmental Disabilities Administration (DDA) enrollment is recommended. DDA is a public agency that provides services and support to qualified individuals. Eligibility differs by state but is typically determined by diagnosis and/or associated cognitive/adaptive disabilities.

- Families with limited income and resources may also qualify for supplemental security income (SSI) for their child with a disability.

## Motor Dysfunction

### Gross motor dysfunction

- Physical therapy is recommended to maximize mobility.
- Consider use of durable medical equipment as needed (e.g., wheelchairs, walkers, bath chairs, orthotics, adaptive strollers).

**Fine motor dysfunction.** Occupational therapy is recommended for difficulty with fine motor skills that affect adaptive function such as feeding, grooming, dressing, and writing.

**Oral motor dysfunction.** Assuming that it is safe for the individual to eat by mouth, feeding therapy, typically from an occupational or speech therapist, is recommended for affected individuals who have difficulty feeding as a result of poor oral motor control.

**Communication issues.** Consider evaluation for alternative means of communication (e.g., [augmentative and alternative communication](#) [AAC]) for individuals who have expressive language difficulties.

## Chapter Notes

### Revision History

- 8 March 2018 (bp) Review posted live
- 13 October 2017 (aeh) Original submission

## References

### Literature Cited

- El-Hattab AW, Craigen WJ, Scaglia F. Mitochondrial DNA maintenance defects. *Biochim Biophys Acta Mol Basis Dis.* 2017;1863:1539–55. PubMed PMID: 28215579.
- Rahbari R, Wuster A, Lindsay SJ, Hardwick RJ, Alexandrov LB, Turki SA, Dominiczak A, Morris A, Porteous D, Smith B, Stratton MR, Hurles ME, et al. Timing, rates and spectra of human germline mutation. *Nat Genet.* 2016;48:126–33. PubMed PMID: 26656846.

## License

GeneReviews® chapters are owned by the University of Washington. Permission is hereby granted to reproduce, distribute, and translate copies of content materials for noncommercial research purposes only, provided that (i) credit for source (<http://www.genereviews.org/>) and copyright (© 1993-2024 University of Washington) are included with each copy; (ii) a link to the original material is provided whenever the material is published elsewhere on the Web; and (iii) reproducers, distributors, and/or translators comply with the [GeneReviews® Copyright Notice and Usage Disclaimer](#). No further modifications are allowed. For clarity, excerpts of GeneReviews chapters for use in lab reports and clinic notes are a permitted use.

For more information, see the [GeneReviews® Copyright Notice and Usage Disclaimer](#).

For questions regarding permissions or whether a specified use is allowed, contact: [admasst@uw.edu](mailto:admasst@uw.edu).