

G.12.1.8 Residential care staff training: awareness and communication

Quality assessment						No of patients		Effect estimate	Quality
No of studies	Design	Risk of bias	Indirectness	Inconsistency	Imprecision	Intervention	Usual care	Summary of results	
Quality of life (QUALID - measured by family member) (higher values favour control)									
1 (Clare 2013)	RCT	Serious ¹	Not serious	N/A	Not serious	32	33	MD -3.98 (-7.60, -0.36)	Moderate
Well-being (Positive Response Schedule) (higher values favour intervention)									
1 (Clare 2013)	RCT	Serious ¹	Not serious	N/A	Serious ²	32	33	MD 2.68 (-3.55, 8.91)	Low
Cognitive function (GADS) (higher values favour control) (higher values favour intervention)									
1 (Clare 2013)	RCT	Serious ¹	Not serious	N/A	Serious ²	32	33	MD -1.18 (-3.44, 1.08)	Low
Behaviour - self-care (Behavioural Assessment Scale of Later Life) (higher values favour intervention)									
1 (Clare 2013)	RCT	Serious ¹	Not serious	N/A	Serious ²	32	33	MD 0.56 (-1.06, 2.18)	Low
Behaviour - sensory abilities (BASOLL) (higher values favour intervention)									
1 (Clare 2013)	RCT	Serious ¹	Not serious	N/A	Serious ²	32	33	MD -0.04 (-0.51, 0.43)	Low
Behaviour - mobility (BASOLL) (higher values favour intervention)									
1 (Clare 2013)	RCT	Serious ¹	Not serious	N/A	Serious ²	32	33	MD -0.18 (-0.47, 0.11)	Low
<ol style="list-style-type: none"> 1. Randomisation by care home, with only a small number of homes in the study 2. Non-significant result 									