

G.12.1.13 Residential care staff training: reducing antipsychotic drug use

Quality assessment						No of patients		Effect estimate	Quality
No of studies	Design	Risk of bias	Indirectness	Inconsistency	Imprecision	Intervention	Usual care	Summary of results	
Proportion taking neuroleptics (lower numbers favour intervention)									
1 (Fossey 2006)	RCT	Serious ¹	N/A	Not serious	Not serious	176	170	RR 0.55 (0.39, 0.76)	Moderate
Fall in past 12 months (lower numbers favour intervention)									
1 (Fossey 2006)	RCT	Serious ¹	N/A	Not serious	Very serious ³	176	170	RR 0.90 (0.59, 1.38)	Very low
Aggression (Cohen-Mansfield agitation score - lower numbers favour intervention)									
1 (Fossey 2006)	RCT	Serious ¹	N/A	Not serious	Serious ²	176	170	MD 0.3 (-8.3, 8.9)	Low
Wellbeing (dementia care mapping - higher numbers favour intervention)									

Quality assessment						No of patients		Effect estimate	Quality
No of studies	Design	Risk of bias	Indirectness	Inconsistency	Imprecision	Intervention	Usual care	Summary of results	
1 (Fossey 2006)	RCT	Serious ¹	N/A	Not serious	Serious ²	176	170	MD -0.2 (-0.5, 0.2)	Low
<ol style="list-style-type: none"> 1. Lack of appropriate blinding 2. Non-significant result 3. 95% CI crosses two lines of a defined MID interval 									