G.12.1.15   Residential care staff training: person-centred showering

Quality assessment						No of patients		Effect estimate	Quality
No of studies	Design	Risk of bias	Indirectness	Inconsistency	Imprecision	Intervention	Usual care	Summary of results	
Any agitation or aggression (%time – higher numbers favour control)									
1 (Sloane 2004)	RCT	Serious[1]	Not serious	N/A	Serious[2]	25	24	MD -8.89 (-23.38, 5.60)	Low
Any physical agitation or aggression (%time – higher numbers favour control)									
1 (Sloane 2004)	RCT	Serious[1]	Not serious	N/A	Serious[2]	25	24	MD -0.39 (-1.67, 0.89)	Low
Any aggression (rate/15minutes – higher numbers favour control)									
1 (Sloane 2004)	RCT	Serious[1]	Not serious	N/A	Not serious	25	24	MD -0.94 (-1.75, -0.13)	Moderate
Hit, bite, kick, throw or spit (rate/15 minutes – higher numbers favour control)									
1 (Sloane 2004)	RCT	Serious[1]	Not serious	N/A	Serious[2]	25	24	MD -0.33 (-0.73, 0.07)	Low

Quality assessment						No of patients		Effect estimate	Quality
No of studies	Design	Risk of bias	Indirectness	Inconsistency	Imprecision	Intervention	Usual care	Summary of results	
Other aggression (attempts/grabbing, rate/15 minutes – higher numbers favour control)									
1 (Sloane 2004)	RCT	Serious[1]	Not serious	N/A	Not serious	25	24	MD -0.78 (-1.54, -0.02)	Moderate
Yelling, crying, moaning (%time – higher numbers favour control)									
1 (Sloane 2004)	RCT	Serious[1]	Not serious	N/A	Serious[2]	25	24	MD -0.09 (-0.69, 0.51)	Low
Complaints, threats (rate/15 minutes – higher numbers favour control)									
1 (Sloane 2004)	RCT	Serious[1]	Not serious	N/A	Serious[2]	25	24	MD -0.39 (-1.35, 0.57)	Low
Mean discomfort scale score (higher numbers favour control)									
1 (Sloane 2004)	RCT	Serious[1]	Not serious	N/A	Not serious	25	24	MD -0.31 (-0.54, -0.08)	Moderate

1. No information on study dropouts
2. Non-significant result