G.12.1.17 Residential care staff training: sensitivity to non-verbal emotion signals

Quality assessment						No of patients		Effect estimate	Quality
No of studies	Design	Risk of bias	Indirectnes	Inconsistenc	Imprecisio	Interventio	Usual	Summary of results	
			S	у	n	n	care		
Symptomatology (higher numbers favour control)									
1 (Magai 2002)	RCT	Very serious ¹	Not serious	N/A	Serious ²	34	23	MD -39.20	Very low
								(-57.15, -21.25)	
Positive emotion (higher numbers favour intervention)									
1 (Magai 2002)	RCT	Very serious ¹	Not serious	N/A	Serious ³	41	27	MD 0.70 (-0.89, 2.29)	Very low
Negative emotion (higher numbers favour control)									
1 (Magai 2002)	RCT	Very serious ¹	Not serious	N/A	Serious ³	41	27	MD 0.10 (-1.49, 1.69)	Very low
Brief symptom inventory (higher numbers favour control)									
1 (Magai 2002)	RCT	Very serious ¹	Not serious	N/A	Serious ³	8	5	MD -4.90	Very low
·								(-14.34, 4.54)	

^{1.} Large differences in baseline characteristics between the intervention and control groups, including in outcome measures

^{2.} Significant differences between the intervention and control groups at baseline in this outcome, which may be a confounding factor in the mean change data

^{3.} Non-significant result

[©] National Institute for Health and Care Excellence, 2017