

## G.12.1.20 Residential care nurse training: managing depression nursing guideline

Quality assessment						No of patients		Effect estimate	Quality
No of studies	Design	Risk of bias	Indirectness	Inconsistency	Imprecision	Intervention	Usual care	Summary of results	
<b>Depression (MDS/RAI-DRS – higher numbers favour control)</b>									
1 (Verkaik 2011)	RCT	Not serious	Not serious	N/A	Serious <sup>1</sup>	62	35	MD -1.00 (-2.41, 0.41)	Moderate
<b>Depression (Cornell Scale – higher numbers favour control)</b>									
1 (Verkaik 2011)	RCT	Not serious	Not serious	N/A	Serious <sup>1</sup>	62	35	MD 0.09 (-2.56, 2.74)	Moderate
<b>Mood (morning care – higher numbers favour intervention)</b>									
1 (Verkaik 2011)	RCT	Not serious	Not serious	N/A	Serious <sup>1</sup>	62	35	MD -0.01 (-0.34, 0.32)	Moderate
<b>Mood (living room – higher numbers favour intervention)</b>									
1 (Verkaik 2011)	RCT	Not serious	Not serious	N/A	Serious <sup>1</sup>	62	35	MD -0.09 (-0.35, 0.17)	Moderate
1. Non-significant result									