

G.12.1.24 GP training: flexible education

Quality assessment						No of patients		Effect estimate	Quality
No of studies	Design	Risk of bias	Indirectness	Inconsistency	Imprecision	Intervention	Usual care	Summary of results	
<b>Quality of life (self-rated) using QOL-AD (higher values favour intervention)</b>									
1 (Beer 2011)	RCT	Not serious	Not serious	N/A	Serious <sup>1</sup>	157	194	MD -0.61 (-3.07, 1.85)	Moderate
<b>Quality of life (carer-rated) using QOL-AD (higher values favour intervention)</b>									
1 (Beer 2011)	RCT	Not serious	Not serious	N/A	Serious <sup>1</sup>	157	194	MD -0.07 (-2.31, 2.17)	Moderate
<b>Quality of life (carer-rated) using ADRQOL (higher values favour intervention)</b>									
1 (Beer 2011)	RCT	Not serious	Not serious	N/A	Serious <sup>1</sup>	157	194	MD 1.02 (-3.23, 5.27)	Moderate
<b>Pain observed (Brief Pain Inventory) (log odds ratio) (higher values favour control)</b>									

Quality assessment						No of patients		Effect estimate	Quality
No of studies	Design	Risk of bias	Indirectness	Inconsistency	Imprecision	Intervention	Usual care	Summary of results	
1 (Beer 2011)	RCT	Not serious	Not serious	N/A	Very serious <sup>2</sup>	157	194	OR 0.60 (0.25, 1.47)	Low
<b>Behavioural and psychological symptoms of dementia (NPI) (higher values favour control)</b>									
1 (Beer 2011)	RCT	Not serious	Not serious	N/A	Very serious <sup>2</sup>	157	194	OR 0.81 (0.40, 1.61)	Low
<b>Use of physical restraint observed (higher values favour control)</b>									
1 (Beer 2011)	RCT	Not serious	Not serious	N/A	Serious <sup>3</sup>	157	194	OR 0.44 (0.17, 1.11)	Moderate
<ol style="list-style-type: none"> <li>1. Non-significant result</li> <li>2. 95% CI crosses two lines of a defined MID</li> <li>3. 95% CI crosses one line of a defined MID</li> </ol>									