

G.3.2 Post diagnosis review for people living with dementia

- How should people living with dementia be reviewed post diagnosis?

G.3.2.1 Managed health services in partnership with Alzheimer’s associations services versus usual managed care services only

Quality assessment						No of patients	Effect estimate	Quality
No of studies	Design	Risk of bias	Indirectness	Inconsistency	Imprecision			
Outcome: Number of emergency department visits at 12 months								
Bass (2003)	RCT	Not serious	Not serious	N/A	Serious ¹	157	MD -0.17 (-0.51, 0.17)	Moderate
Outcome: Number of hospital admissions at 12 months								
Bass (2003)	RCT	Not serious	Not serious	N/A	Serious ¹	157	MD -0.08 (-0.26, 0.10)	Moderate
Outcome: Number of physician visits at 12 months								
Bass (2003)	RCT	Not serious	Not serious	N/A	Serious ¹	157	MD 0.01 (-1.36, 1.38)	Moderate
Outcome: Use of case management at 12 months								
Bass (2003)	RCT	Not serious	Not serious	N/A	Not serious	157	MD -0.16 (-0.29, -0.03)	High
Outcome: Use of direct care community services at 12 months								
Bass (2003)	RCT	Not serious	Not serious	N/A	Serious ¹	157	MD 0.02 (-0.47, 0.51)	Moderate
Outcome: Use of non-Association information and support services								
Bass (2003)	RCT	Not serious	Not serious	N/A	Serious ¹	157	MD -0.10 (-0.50, 0.30)	Moderate
1. Non-significant result								