

**G.4.1.2 Family-centred function focused care versus usual care**

Quality assessment						No of patients		Effect estimate	Quality
No of studies	Design	Risk of bias	Indirectness	Inconsistency	Imprecision	Intervention	Comparator		
<b>Outcome: Mean difference in length of stay at discharge</b>									
Boltz (2015)	Non randomised controlled trial	Very serious <sup>1</sup>	Not serious	N/A	Serious <sup>2</sup>	44	42	MD -0.40 (-1.27, 0.47)	Very low
<b>Outcome: Hospital readmissions at 30 days</b>									
Boltz (2015)	Non randomised controlled trial	Very serious <sup>1</sup>	Not serious	N/A	Serious <sup>2</sup>	44	42	MD -7.00 (-14.55, 0.55)	Very low
<b>Outcome: Utilisation of post-acute rehabilitation at discharge</b>									
Boltz (2015)	Non randomised controlled trial	Very serious <sup>1</sup>	Not serious	N/A	Serious <sup>2</sup>	44	42	MD 2.00 (-25.48, 29.48)	Very low
<b>Outcome: Activities of Daily Living (Barthel Index) at 2 months</b>									
Boltz (2015)	Non randomised controlled trial	Very serious <sup>1</sup>	Not serious	N/A	Not serious	44	42	MD 20.7 (10.32, 31.08)	Low
<b>Outcome: Walking performance (50 yards) at 2 months</b>									
Boltz (2015)	Non randomised controlled trial	Very serious <sup>1</sup>	Not serious	N/A	Not serious	44	42	MD 5.60 (3.39, 7.81)	Low
<b>Outcome: Gait and Balance (Tinetti Scale) at 2 months</b>									

Quality assessment						No of patients		Effect estimate	Quality
No of studies	Design	Risk of bias	Indirectness	Inconsistency	Imprecision	Intervention	Comparator		
Boltz (2015)	Non randomised controlled trial	Very serious <sup>1</sup>	Not serious	N/A	Serious <sup>2</sup>	44	42	MD 1.50 (-2.39, 5.39)	Very low
<b>Outcome: Delirium severity (Delirium severity Scale) at 2 months</b>									
Boltz (2015)	Non randomised controlled trial	Very serious <sup>1</sup>	Not serious	N/A	Not serious	44	42	MD -2.00 (-3.09, -0.91)	Low
<b>Outcome: Delirium present at 2 months post discharge</b>									
Boltz (2015)	Non randomised controlled trial	Very serious <sup>1</sup>	Not serious	N/A	Not serious	44	42	MD -9.00 (-17.83, -0.17)	Low
<b>Outcome: Carer preparedness for caregiving at 2 months</b>									
Boltz (2015)	Non randomised controlled trial	Very serious <sup>1</sup>	Not serious	N/A	Serious <sup>2</sup>	44	42	MD -3.10 (-5.73, 0.47)	Very low
<b>Outcome: Carer anxiety (HADS-A) at 2 months</b>									
Boltz (2015)	Non randomised controlled trial	Very serious <sup>1</sup>	Not serious	N/A	Serious <sup>2</sup>	44	42	MD -1.60 (-3.57, 0.37)	Very low
<b>Outcome: Carer depression (HADS-D) at 2 months</b>									
Boltz (2015)	Non randomised controlled trial	Very serious <sup>1</sup>	Not serious	N/A	Serious <sup>2</sup>	44	42	MD -0.70 (-2.54, 1.14)	Very low
<b>Outcome: Carer role strain (Modified Caregiver Strain Index) at 2 months</b>									
Boltz (2015)	Non randomised controlled trial	Very serious <sup>1</sup>	Not serious	N/A	Serious <sup>2</sup>	44	42	MD -0.80 (-3.06, 1.46)	Very low
<b>Outcome: Carer mutuality at 2 months</b>									
Boltz (2015)	Non randomised controlled trial	Very serious <sup>1</sup>	Not serious	N/A	Serious <sup>2</sup>	44	42	MD 3.50 (-1.51, 8.51)	Very low
<ol style="list-style-type: none"> <li>1. Non-randomised study; high risk of bias based on limited reporting of study.</li> <li>2. Non-significant result.</li> </ol>									