G.11.1.3 Psychoeducation and skills training

Quality assessment						No of patients		Effect estimate	Quality		
No of studies	Design	Risk of bias	Indirectness	Inconsistency	Imprecision	Intervention	Usual care	Summary of results			
Carer burden (lower values favour intervention)											
10	RCT	Not serious	Not serious	Serious ¹	Serious ²	595	551	SMD -0.30 (-0.49, -0.10)	Low		
Carer depression (lower values favour intervention)											
14	RCT	Not serious	Not serious	Not serious	Serious ²	1,102	929	SMD -0.25 (-0.33, -0.16)	Moderate		
Carer anxiety (lower values favour intervention)											
6	RCT	Not serious	Not serious	Not serious	Serious ²	606	483	SMD -0.26 (-0.39, -0.14)	Moderate		
Carer stress (lower values favour intervention)											
6	RCT	Not serious	Not serious	Not serious	Serious ²	323	323	SMD -0.21 (-0.37, -0.06)	Moderate		
Carer quality of life (higher values favour intervention)											
5	RCT	Not serious	Not serious	Serious ¹	Serious ²	334	324	SMD 0.11 (-0.11, 0.33)	Low		
Carer self-efficacy (higher values favour intervention)											
7	RCT	Not serious	Not serious	Serious ¹	Serious ²	503	470	SMD 0.20 (-0.01, 0.42)	Low		
Revised memory and behaviour problems checklist – severity (lower values favour intervention)											
3	RCT	Not serious	Not serious	Serious ¹	Very serious ³	115	92	SMD -0.11 (-0.52, 0.30)	Very low		
Revised memory and behaviour problems checklist – reaction (lower values favour intervention)											
2	RCT	Not serious	Not serious	Serious ¹	Serious ²	211	172	SMD -0.24 (-0.54, 0.07)	Low		
Behavioural and psychological symptoms of dementia – severity (lower values favour intervention)											

[©] National Institute for Health and Care Excellence, 2017

Quality assessment							atients	Effect estimate	Quality		
No of studies	Design	Risk of bias	Indirectness	Inconsistency	Imprecision	Intervention	Usual care	Summary of results			
7	RCT	Not serious	Not serious	Serious ¹	Serious ²	295	289	SMD -0.27 (-0.53, -0.02)	Low		
Behavioural and psychological symptoms of dementia – reaction (lower values favour intervention)											
3	RCT	Not serious	Not serious	Not serious	Serious ²	68	74	SMD -0.23 (-0.56, 0.10)	Moderate		
Activities of daily living – person living with dementia (higher values favour intervention)											
3	RCT	Not serious	Not serious	Not serious	Serious ²	128	133	SMD -0.07 (-0.31, 0.18)	Moderate		
Proportion entering long stay care (lower values favour intervention)											
3	RCT	Not serious	Not serious	Not serious	Serious ²	265	195	RR 1.47 (0.91, 2.37)	Moderate		
 i²>40% Crosses one line of a defined MID Crosses two lines of a defined MID 											

[©] National Institute for Health and Care Excellence, 2017