G.11.1.8 Multicomponent interventions

Quality assessment						No of patients		Effect estimate	Quality
No of studies	Design	Risk of bias	Indirectness	Inconsistency	Imprecision	Intervention	Usual care	Summary of results	
Carer burden (lo	wer values	favour intervention	1)						
15	RCT	Not serious	Not serious	Serious ¹	Serious ²	1,663	1,581	SMD -0.17 (-0.33, -0.01)	Low
Carer depressior	ı (lower val	ues favour interve	ntion)						
20	RCT	Not serious	Not serious	Serious ¹	Serious ²	2,806	2,414	SMD -0.29 (-0.49, -0.09)	Low
Carer anxiety (lo	wer values	favour intervention	n)						
2	RCT	Not serious	Not serious	Not serious	Very serious ³	43	35	SMD 0.05 (-0.40, 0.50)	Low
Carer quality of I	ife (higher v	alues favour inter	vention)						
3	RCT	Not serious	Not serious	Serious ¹	Serious ²	337	343	SMD 0.34 (0.04, 0.64)	Low
Carer self-efficac	y (higher v	alues favour interv	vention)						
1 (Martin-Cook 2005)	RCT	Not serious	Not serious	N/A	Very serious ³	24	23	SMD 0.24 (-0.34, 0.81)	Low
Carer social sup	port (higher	values favour inte	ervention)						
2	RCT	Not serious	Not serious	Not serious	Not serious	60	62	SMD 0.56 (0.20, 0.92)	High
Revised memory	and behav	iour problems che	cklist – severity	(lower values favo	our intervention)				
4	RCT	Not serious	Not serious	Not serious	Serious ²	805	549	SMD -0.12 (-0.23, -0.01)	Moderate
Revised memory	and behav	iour problems che	cklist - reaction	(lower values favo	our intervention)				
4	RCT	Not serious	Not serious	Not serious	Serious ²	282	272	SMD -0.19 (-0.43, 0.06)	Moderate

[©] National Institute for Health and Care Excellence, 2017

Quality assessment						No of patients		Effect estimate	Quality		
No of studies	Design	Risk of bias	Indirectness	Inconsistency	Imprecision	Intervention	Usual care	Summary of results			
Behavioural and psychological symptoms of dementia – severity (lower values favour intervention)											
8	RCT	Not serious	Not serious	Serious ¹	Serious ²	465	479	SMD -0.29 (-0.64, 0.07)	Low		
Behavioural and psychological symptoms of dementia – reaction (lower values favour intervention)											
6	RCT	Not serious	Not serious	Not serious	Serious ²	391	409	SMD -0.31 (-0.45, -0.18)	Moderate		
Activities of daily living – person living with dementia (higher values favour intervention)											
6	RCT	Not serious	Not serious	Serious ¹	Serious ²	430	455	SMD 0.33 (-0.15, 0.81)	Low		
Proportion entering long stay care (lower values favour intervention)											
7	RCT	Not serious	Not serious	Serious ¹	Serious ²	520	472	RR 0.80 (0.61, 1.04)	Low		
 i²>40% Crosses one line of a defined MID 											

^{3.} Crosses two lines of a defined MID