

P.2.7.13 Amyloid Beta 1-42/1- 40

| Studies | Design | Total N | Sens (95%CI) | Spec (95%CI) | Measure | Summary of findings (95%CI) | Risk of bias | Inconsistency | Indirectness | Imprecision | Other considerations | Quality |
|---|-----------------|---------|-------------------|-------------------|---------|-----------------------------|--------------|---------------|--------------|-------------|----------------------|----------|
| SECONDARY CARE | | | | | | | | | | | | |
| 3 studies (Dumurgier 2015 (Lille); Dumurgier 2015 (Paris); Dumurgier 2015 (Montpellier)) | 3 × prospective | 367 | 0.83 (0.60, 0.94) | 0.77 (0.66, 0.85) | LR+ | 3.33 (2.31, 4.78) | V. serious | Not serious | Not serious | Not serious | - | LOW |
| | | | | | LR- | 0.22 (0.09, 0.54) | V. serious | Serious | Not serious | Serious | | VERY LOW |
| <p>Notes on risk of bias</p> <p>Dumurgier 2015: The reference standard diagnosis included consideration of the CSF results; the test cut offs were not pre-specified; patients with unknown clinical diagnoses or MCI were excluded from the study and the timing of the reference and index tests is unclear.</p> <p>Additional notes: the Dumurgier study had 3 independent data sets from 3 different clinics.</p> | | | | | | | | | | | | |