

Table 9: Review protocol: Hip replacement surgery

| ID | Field | Content |
|----|------------------------------|---|
| 0. | PROSPERO registration number | Not registered |
| 1. | Review title | Surgical approaches for hip replacement surgery |
| 2. | Review question | In adults having primary elective hip replacement, what is the most clinical and cost-effective approach: posterior, direct anterior, anterolateral, direct superior or SuperPATH? |
| 3. | Objective | There are a number of surgical approaches for hip replacement that can be used. They vary in terms of how invasive the surgery is, the surgeon's access to and visibility of the joint, recovery period after the surgery and limitations in terms of movement and risks of adverse events during or after the surgery. Where multiple approaches are possible there is currently variation in practice and this review seeks to find the most clinically and cost effective approach where there are no contraindications. |
| 4. | Searches | <p>The following databases will be searched: Cochrane Central Register of Controlled Trials (CENTRAL) Cochrane Database of Systematic Reviews (CDSR) Embase MEDLINE</p> <p>Searches will be restricted by: English language Human studies Letters and comments are excluded.</p> <p>Other searches: Inclusion lists of relevant systematic reviews will be checked by the reviewer.</p> <p>The searches may be re-run 6 weeks before final committee meeting and further studies retrieved for inclusion if relevant.</p> |

| ID | Field | Content |
|-----|---|---|
| | | The full search strategies will be published in the final review. |
| 5. | Condition or domain being studied | Hip replacement surgery |
| 6. | Population | <p>Inclusion: Adults indicated for primary total hip replacement</p> <p>Exclusion: Adults having joint replacement as immediate treatment following fracture. Adults having revision joint replacement. Adults having joint replacement as treatment for primary or secondary cancer affecting the bones.</p> |
| 7. | Intervention/Exposure/T est | <p>Primary total hip replacement utilising the posterior approach</p> <p>Primary total hip replacement utilising the direct anterior approach</p> <p>Primary total hip replacement utilising the anterolateral approach</p> <p>Primary total hip replacement utilising the SuperPATH approach</p> <p>Primary total hip replacement utilising the direct superior approach</p> |
| 8. | Comparator/Reference standard/Confounding factors | Comparison between interventions |
| 9. | Types of study to be included | <p>Randomised controlled trials</p> <p>If no well conducted RCTs are available then observational studies with multivariate analysis will be investigated.</p> |
| 10. | Other exclusion criteria | <p>Non-English language studies.</p> <p>Abstracts will be excluded as it is expected there will be sufficient full text published studies available.</p> |
| 11. | Context | N/A |
| 12. | Primary outcomes (critical outcomes) | <p>Mortality: life expectancy (dichotomous)</p> <p>Mortality: 30 day (dichotomous)</p> <p>Quality of life at 6 weeks or earlier, later than 6 weeks up to 1 year, at least 2 years (continuous)</p> <p>Patient Reported Outcome Measures (PROMs) at 6 weeks or earlier, later than 6 weeks up to 1 year, at least 2 years (continuous)</p> |

| ID | Field | Content |
|-----|---|---|
| | | Revision rate of joint replacement (time to event) |
| 13. | Secondary outcomes (important outcomes) | <p>Deep surgical site infection (dichotomous) Superficial surgical site infection (dichotomous) Length of stay (continuous) Reoperation/dislocation rate (dichotomous) Intraoperative complications (for example nerve damage) Surgery time (continuous)</p> <p>To be extracted when not included within a PROM: Function at 6 weeks or earlier, later than 6 weeks up to 1 year, at least 2 years (continuous) Pain at 6 weeks or earlier, later than 6 weeks up to 1 year, at least 2 years (continuous)</p> |
| 14. | Data extraction (selection and coding) | <p>EndNote will be used for reference management, sifting, citations and bibliographies. Titles and/or abstracts of studies retrieved using the search strategy and those from additional sources will be screened for inclusion. The full text of potentially eligible studies will be retrieved and will be assessed for eligibility in line with the criteria outlined above.</p> <p>10% of the abstracts will be reviewed by two reviewers, with any disagreements resolved by discussion or, if necessary, a third independent reviewer.</p> <p>An in-house developed database; EviBase, will be used for data extraction. A standardised form is followed to extract data from studies (see Developing NICE guidelines: the manual section 6.4) and for undertaking assessment of study quality. Summary evidence tables will be produced including information on: study setting; study population and participant demographics and baseline characteristics; details of the intervention and control interventions; study methodology' recruitment and missing data rates; outcomes and times of measurement; critical appraisal ratings.</p> <p>A second reviewer will quality assure the extracted data. Discrepancies will be identified and resolved through discussion (with a third reviewer where necessary).</p> |
| 15. | Risk of bias (quality) assessment | <p>Risk of bias will be assessed using the appropriate checklist as described in Developing NICE guidelines: the manual. For Intervention reviews the following checklist will be used according to study design being assessed: Systematic reviews: Risk of Bias in Systematic Reviews (ROBIS) Randomised Controlled Trial: Cochrane RoB (2.0)</p> <p>Disagreements between the review authors over the risk of bias in particular studies will be resolved by discussion, with involvement of a third review author where necessary.</p> |
| 16. | Strategy for data | Where possible, data will be meta-analysed. Pairwise meta-analyses will be performed using Cochrane Review Manager |

| ID | Field | Content | | | | | | | | | | | | | | |
|-------------------------------------|---------------------------|---|-------------------------------------|--------------|--------------------------|------------|--------------------------|------------|--------------------------|-------------|--------------------------|---------------|--------------------------|------------------|--------------------------|------------------------|
| | synthesis | <p>(RevMan5) to combine the data given in all studies for each of the outcomes stated above. A fixed effect meta-analysis, with weighted mean differences for continuous outcomes and risk ratios for binary outcomes will be used, and 95% confidence intervals will be calculated for each outcome.</p> <p>Heterogeneity between the studies in effect measures will be assessed using the I² statistic and visually inspected. We will consider an I² value greater than 50% indicative of substantial heterogeneity. Sensitivity analyses will be conducted based on pre-specified subgroups using stratified meta-analysis to explore the heterogeneity in effect estimates. If this does not explain the heterogeneity, the results will be presented using random-effects.</p> <p>GRADE pro will be used to assess the quality of each outcome, taking into account individual study quality and the meta-analysis results. The 4 main quality elements (risk of bias, indirectness, inconsistency and imprecision) will be appraised for each outcome.</p> <p>If the population included in an individual study includes children aged under 12, it will be included if the majority of the population is aged over 12, and downgraded for indirectness if the overlap into those aged less than 12 is greater than 20%.</p> <p>Publication bias is tested for when there are more than 5 studies for an outcome. Other bias will only be taken into consideration in the quality assessment if it is apparent.</p> <p>Where meta-analysis is not possible, data will be presented and quality assessed individually per outcome.</p> <p>If sufficient data is available to make a network of treatments, WinBUGS will be used for network meta-analysis.</p> | | | | | | | | | | | | | | |
| 17. | Analysis of sub-groups | <p>Age: Working age Non-working age</p> | | | | | | | | | | | | | | |
| 18. | Type and method of review | <table border="1"> <tr> <td data-bbox="678 1142 1361 1174"><input checked="" type="checkbox"/></td> <td data-bbox="1373 1142 2110 1174">Intervention</td> </tr> <tr> <td data-bbox="678 1182 1361 1214"><input type="checkbox"/></td> <td data-bbox="1373 1182 2110 1214">Diagnostic</td> </tr> <tr> <td data-bbox="678 1222 1361 1254"><input type="checkbox"/></td> <td data-bbox="1373 1222 2110 1254">Prognostic</td> </tr> <tr> <td data-bbox="678 1262 1361 1294"><input type="checkbox"/></td> <td data-bbox="1373 1262 2110 1294">Qualitative</td> </tr> <tr> <td data-bbox="678 1302 1361 1334"><input type="checkbox"/></td> <td data-bbox="1373 1302 2110 1334">Epidemiologic</td> </tr> <tr> <td data-bbox="678 1342 1361 1374"><input type="checkbox"/></td> <td data-bbox="1373 1342 2110 1374">Service Delivery</td> </tr> <tr> <td data-bbox="678 1382 1361 1414"><input type="checkbox"/></td> <td data-bbox="1373 1382 2110 1414">Other (please specify)</td> </tr> </table> | <input checked="" type="checkbox"/> | Intervention | <input type="checkbox"/> | Diagnostic | <input type="checkbox"/> | Prognostic | <input type="checkbox"/> | Qualitative | <input type="checkbox"/> | Epidemiologic | <input type="checkbox"/> | Service Delivery | <input type="checkbox"/> | Other (please specify) |
| <input checked="" type="checkbox"/> | Intervention | | | | | | | | | | | | | | | |
| <input type="checkbox"/> | Diagnostic | | | | | | | | | | | | | | | |
| <input type="checkbox"/> | Prognostic | | | | | | | | | | | | | | | |
| <input type="checkbox"/> | Qualitative | | | | | | | | | | | | | | | |
| <input type="checkbox"/> | Epidemiologic | | | | | | | | | | | | | | | |
| <input type="checkbox"/> | Service Delivery | | | | | | | | | | | | | | | |
| <input type="checkbox"/> | Other (please specify) | | | | | | | | | | | | | | | |

| ID | Field | Content | | |
|-----|--|--|--------------------------|-------------------------------------|
| 19. | Language | English | | |
| 20. | Country | England | | |
| 21. | Anticipated or actual start date | 12/02/19 | | |
| 22. | Anticipated completion date | 31/07/19 | | |
| 23. | Stage of review at time of this submission | Review stage | Started | Completed |
| | | Preliminary searches | <input type="checkbox"/> | <input checked="" type="checkbox"/> |
| | | Piloting of the study selection process | <input type="checkbox"/> | <input checked="" type="checkbox"/> |
| | | Formal screening of search results against eligibility criteria | <input type="checkbox"/> | <input checked="" type="checkbox"/> |
| | | Data extraction | <input type="checkbox"/> | <input checked="" type="checkbox"/> |
| | | Risk of bias (quality) assessment | <input type="checkbox"/> | <input checked="" type="checkbox"/> |
| | | Data analysis | <input type="checkbox"/> | <input checked="" type="checkbox"/> |
| 24. | Named contact | 5a. Named contact National Guideline Centre 5b Named contact e-mail 5e Organisational affiliation of the review National Institute for Health and Care Excellence (NICE) and the National Guideline Centre | | |
| 25. | Review team members | From the National Guideline Centre: Mr Carlos Sharpin [Guideline lead] Mr Alex Allen [Senior Systematic Reviewer] Ms Rafina Yarde [Systematic reviewer] Mr Robert King [Health economist] Ms Agnès Cuyàs [Information specialist] | | |

| ID | Field | Content | |
|-----|--|---|--|
| | | Ms Eleanor Priestnall [Project Manager] | |
| 26. | Funding sources/sponsor | This systematic review is being completed by the National Guideline Centre which receives funding from NICE. | |
| 27. | Conflicts of interest | All guideline committee members and anyone who has direct input into NICE guidelines (including the evidence review team and expert witnesses) must declare any potential conflicts of interest in line with NICE's code of practice for declaring and dealing with conflicts of interest. Any relevant interests, or changes to interests, will also be declared publicly at the start of each guideline committee meeting. Before each meeting, any potential conflicts of interest will be considered by the guideline committee Chair and a senior member of the development team. Any decisions to exclude a person from all or part of a meeting will be documented. Any changes to a member's declaration of interests will be recorded in the minutes of the meeting. Declarations of interests will be published with the final guideline. | |
| 28. | Collaborators | Development of this systematic review will be overseen by an advisory committee who will use the review to inform the development of evidence-based recommendations in line with section 3 of Developing NICE guidelines: the manual. Members of the guideline committee are available on the NICE website: [NICE guideline webpage]. | |
| 29. | Other registration details | | |
| 30. | Reference/URL for published protocol | | |
| 31. | Dissemination plans | NICE may use a range of different methods to raise awareness of the guideline. These include standard approaches such as: notifying registered stakeholders of publication publicising the guideline through NICE's newsletter and alerts issuing a press release or briefing as appropriate, posting news articles on the NICE website, using social media channels, and publicising the guideline within NICE. | |
| 32. | Keywords | hip replacement surgery, approach, hip arthroplasty. | |
| 33. | Details of existing review of same topic by same authors | N/A | |
| 34. | Current review status | <input type="checkbox"/> | Ongoing |
| | | <input checked="" type="checkbox"/> | Completed but not published |
| | | <input type="checkbox"/> | Completed and published |
| | | <input type="checkbox"/> | Completed, published and being updated |

| ID | Field | Content | |
|-----|------------------------------|--|--------------|
| | | <input type="checkbox"/> | Discontinued |
| 35. | Additional information | N/A | |
| 36. | Details of final publication | www.nice.org.uk | |