

E.2 Antidepressants (duloxetine, fluoxetine, moclobemide) versus placebo

Figure 21: Quality of life: SF-36 subscales at 12 weeks (duloxetine)





Figure 22: Fatigue: 14-item Chalder fatigue scale at 26 weeks (fluoxetine)


Figure 23: Fatigue: MFI-20 at 12 weeks (duloxetine)




Figure 24: Fatigue: Checklist Individual Strength (CIS) fatigue at 16 weeks (fluoxetine)**Figure 25: Physical functioning: Karnofsky Performance Index at 6 weeks (moclobemide)**

Figure 26: Psychological status: Profile of mood states (POMS) at 6 weeks (moclobemide)Footnotes

(1) Vigour subscale inverted for analysis




Figure 27: Psychological status: HADS depression at 12-26 weeks (change scores) (fluoxetine or duloxetine)**Figure 28: Psychological status: HADS anxiety at 12 weeks (duloxetine)****Figure 29: Psychological status: Beck Depression Inventory at 16 weeks (fluoxetine)**


Figure 30: Pain: Brief Pain Inventory at 12 weeks (duloxetine)



Figure 31: Adverse events (fluoxetine)**Figure 32: Exercise performance measure: VO₂ max (mL O₂/kg/min) at 26 weeks (fluoxetine)**




Figure 33: Symptom scales: Clinical Global Impression at 12 weeks (duloxetine)**Figure 34: Symptom scales: CDC symptom inventory at 12 weeks (duloxetine)**

Figure 35: Symptom scales: Improvement of symptoms (patient-reported) at 6-14 weeks (fluoxetine or moclobemide)**Figure 36: Symptom scales: Worsening of symptoms (patient-reported) at 14 weeks (fluoxetine)**